

Quantum Breathing Mode for Electrons with $1/r^2$ Interaction

Michael R. Geller and Giovanni Vignale

Department of Physics, University of Missouri, Columbia, Missouri, 65211

(February 6, 2008)

In a recent Letter [1], Johnson and Quiroga have obtained several interesting exact results for electrons with $1/r^2$ interactions in a two-dimensional quantum dot. A parabolic confining potential of the form $\frac{1}{2}m\omega_0^2 r^2$ is assumed, and the system is subjected to a uniform perpendicular magnetic field of strength B . In particular, they have shown that there exists a collective “breathing” mode excitation with frequency

$$\omega = 2\Omega, \quad (1)$$

where $\Omega^2 \equiv \omega_0^2 + \frac{1}{4}\omega_c^2$, $\omega_c \equiv eB/mc$ is the cyclotron frequency, and m is the effective mass. The exact spectrum of the interacting electron system therefore contains an infinite ladder of energy levels at integer multiples of $2\hbar\Omega$.

The purpose of this Comment is to point out that the quantum breathing mode excitation spectrum can also be obtained directly from the property of the many-particle Hamiltonian under a certain scale transformation, and in a manner that makes evident the special property of the inverse-square-law interaction for the quantum breathing mode. We shall work in the symmetric gauge and write the Hamiltonian as $H = T + \frac{1}{2}\omega_c L_z + V + U$, where $T \equiv \sum_n p_n^2/2m$, with $\mathbf{p}_n$ the canonical momentum, $L_z \equiv \sum_n (\mathbf{r}_n \times \mathbf{p}_n) \cdot \hat{z}$ is the z component of the canonical angular momentum, $V \equiv \sum_n \frac{1}{2}m\Omega^2 r_n^2$ is the effective field-dependent parabolic confining potential, and

$$U \equiv \sum_{n < n'} \frac{g}{|\mathbf{r}_n - \mathbf{r}_{n'}|^\alpha} \quad (2)$$

is any *power-law* electron-electron interaction.

We first note the properties of H under a scale transformation $O \rightarrow e^{i\lambda S} O e^{-i\lambda S}$ generated by

$$S \equiv \frac{1}{2} \sum_n (\mathbf{r}_n \cdot \mathbf{p}_n + \mathbf{p}_n \cdot \mathbf{r}_n). \quad (3)$$

This transformation performs a radial displacement of each coordinate by an amount proportional to its distance from the origin; that is, it generates a “breathing” motion. Under this transformation, $T \rightarrow T - 2\lambda T$, $L_z \rightarrow L_z$, $V \rightarrow V + 2\lambda V$, and $U \rightarrow U - \alpha\lambda U$, to first order in λ . Therefore,

$$H \rightarrow H + i\lambda[S, H] = H - 2\lambda T + 2\lambda V - \alpha\lambda U. \quad (4)$$

Equation (4), in turn, may be regarded as an equation of motion for S . In fact, noting that $dV/dt = \Omega^2 S$, we obtain the operator equation of motion

$$\frac{d^2 V}{dt^2} + (2 + \alpha)\Omega^2 V = \alpha\Omega^2 (H - \frac{1}{2}\omega_c L_z) + (2 - \alpha)\Omega^2 T, \quad (5)$$

which is the same as one would obtain classically.

The breathing mode of the corresponding classical system of point charges may be obtained from (5) by considering small oscillations about an equilibrium configuration, where the velocities are zero. Because H and L_z are constants of the motion, whereas the physical kinetic energy $T + \frac{1}{2}\omega_c L_z + \omega_c^2 V/4\Omega^2$ is zero to first order in the displacements, the *classical* breathing mode frequency is generally $\omega = \sqrt{(2 + \alpha)\omega_0^2 + \omega_c^2}$. For example, the classical breathing frequency of electrons with Coulomb interactions ($\alpha = 1$) in a parabolic dot with no magnetic field is $\sqrt{3}\omega_0$, a result first obtained by Schweigert and Peeters [2].

Quantum zero-point motion, however, generally modifies the breathing mode and the other classical normal modes, by shifting their frequencies and by giving them a finite lifetime of the order of a/R , with a denoting the Bohr radius and R the radius of the droplet of charge in the dot. An exception occurs when $\alpha = 2$: In this case V becomes an *exact* quantum collective coordinate with frequency (1), independent of g and N . The collective coordinate V may be separated into a center-of-mass and relative-coordinate part, $V = V_{\text{cm}} + V_{\text{rel}}$. For $\alpha = 2$, it can be shown that each component separately satisfies a harmonic oscillator equation of motion of the form (5) with frequency 2Ω . V_{rel} is the collective coordinate corresponding to the breathing mode discovered by Johnson and Quiroga [1].

This work was supported by the NSF Grants No. DMR-9403908 and DMR-9416906. We gratefully acknowledge the hospitality of the Condensed Matter Theory Group at Indiana University, where this work was initiated, and we thank Allan MacDonald for stimulating discussions, and Francois Peeters for first drawing our attention to the breathing mode in classical systems.

[1] N. F. Johnson and L. Quiroga, Phys. Rev. Lett. **74**, 4277 (1995).

[2] V. A. Schweigert and F. M. Peeters, Phys. Rev. B **51**, 7700 (1995); F. M. Peeters, V. A. Schweigert and V. M. Bedanov, Physica B **212**, 710 (1995).